

**HARMFUL**  
**KEEP OUT OF REACH OF CHILDREN**

**RODENTHOR**<sup>®</sup>  
**SOFT BAIT RODENTICIDE**

**Active ingredient: Contains 0.05 g/kg brodifacoum  
in the form of a ready to use block bait.**

For use in damp or dry situations in  
and around industrial, commercial, public  
services, agricultural and domestic buildings,  
including food handling establishments.  
Controls rodents resistant to other anticoagulants,  
including warfarin, coumatetralyl and bromadiolone.

**ONE FEED KILLS RATS AND MICE\***

**REGISTERED TO & MANUFACTURED BY:**

Zapl SpA  
Via Terza Strada 12, 35026 Conselve, Padua, Italy  
Registered pursuant to the ACVM Act 1997, No. V9579  
See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.

Approved pursuant to the HSNO Act 1996,  
Approval Code: HSR100814

See [www.epa.govt.nz](http://www.epa.govt.nz) for approval controls.

**NZ AGENT:**

ENSYSTEX NEW ZEALAND LTD  
17C Corinthian Drive, Albany 0752, Auckland  
[www.Ensystex.co.nz](http://www.Ensystex.co.nz)  
CUSTOMER SERVICE: 0800 ENSYSTEX (0800 367 978)

**Net Contents:**

**9 kg**

© Registered trademarks of Ensystex, Inc. used by licence to Ensystex New Zealand Ltd.

**ENSYSTEX**<sup>®</sup>



BY LAW THIS PRODUCT MUST ONLY BE USED AS SPECIFIED ON THIS LABEL.

HSNO Classifications: 6.1E, 6.5B, 6.9B, 9.1D, 9.3B.

## TOXICITY

**Harmful – This product may be harmful if swallowed.**

**May cause sensitisation from prolonged skin contact and may cause organ damage from repeated oral exposure at high doses.**

## FIRST AID

If swallowed do not induce vomiting.

For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. Vitamin K1 (Phytomenadione) is antidotal.

*Note to Physicians* - brodifacoum is a long lasting anticoagulant chemical, which if ingested by humans, domestic animals or pets, can reduce the clotting power of the blood and haemorrhage may result. Vitamin K1 (Phytomenadione) should be administered urgently by medical or appropriate para-medical personnel. If administered intravenously the injection must be given slowly.

## DIRECTIONS FOR USE

### Restraints

Avoid contact with food, food utensils, or places where food is prepared or stored.

### Baiting Strategy

- Eliminate as far as practical, all alternative food sources.
- Bait infested area. Place baits under cover.
- Actively search for, remove and destroy dead and moribund rodents whilst bait is being used.
- Ensure baits are removed when rodent activity has ceased.
- Always record the number and locations of baits for the purpose of bait replacement and later removal.
- Nails or wire may be used to fix baits to prevent them being removed by rodents. Alternatively, baits may be secured in bait stations.
- Place baits so as to prevent access to non-target animals. Use tamper-resistant bait stations at locations where access by non-target animals cannot otherwise be prevented, and fix the stations securely to the substrate where necessary.
- Use a greater number of baits and shorter distance between bait placements when infestations are heavy.
- Conduct initial inspection of bait points after 3 or 4 days. Replace any bait that has been consumed. Repeat approximately 4 days later and then weekly. Replace contaminated or spoiled baits.

## ECOTOXICITY

**Toxic to rodents and other terrestrial vertebrates.**

**RODENTHOR SOFT BAIT Rodenticide must not be applied onto or into water.**

**RODENTHOR SOFT BAIT Rodenticide must be applied as a contained, ground-based application only.**

- Baiting for at least 2 weeks (3 to 4 pulses) may be necessary to reduce rat/mouse numbers to a low level. Although heavy infestations may require longer treatments to achieve complete eradication.
- Discontinue the treatment when effective control has been achieved and remove all bait and bait stations unless a maintenance treatment is to be undertaken

## Application

Situation	Pests	Rate	Critical Comments
Industrial, Commercial, Public Services, Agricultural and Domestic Buildings, including food handling establishments	Rats ( <i>Rattus rattus</i> , <i>R. norvegicus</i> ), Mice ( <i>Mus musculus</i> )	1 to 3 blocks/ bait station	Set out bait in places frequented by mice and rats e.g. in and around buildings, fence lines, enclosed spaces or drains. <b>Clean-Out Treatment:</b> <b>For rats:</b> Place 1 to 3 sachets at intervals of 5 to 9 m in infested areas. <b>For mice:</b> Place single sachets at 2 to 3 m intervals in infested areas.

## GENERAL INSTRUCTIONS

Most deaths occur 4 to 7 days after consumption of bait. Locate sachets along walls or rafters and in the dark, sheltered spots where there are signs of rodent activity.

\*Note: One feed is sufficient to kill rats and mice. Although rats and mice will continue to eat the bait after a lethal dose has been eaten, the directions for use are intended to minimize this.

Depending on the amount eaten, this bait may kill or injure. The bait is hazardous to dogs, cats, pigs, poultry and other wildlife. Do NOT contaminate dams, waterways and drains with the bait or its used container.

## STORAGE

Store in the closed original container in a cool, well ventilated area. Do NOT store for prolonged periods in direct sunlight. Do NOT store near food or feedstuffs.

## PRECAUTIONS

Keep out of reach of children.

Do NOT inhale dust or touch bait sachets with bare hands.

Do NOT place bait sachets in the open or places accessible to children, domestic pets, animals or birds. After use, wash hands and exposed skin thoroughly with soap and water

## DISPOSAL

Break, crush, puncture and bury empty containers and dispose of unused bait in a local authority landfill. If no landfill is available, bury the containers or unused bait below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. Do NOT use containers which have held bait for any other purpose. Dispose of carcasses safely by burning or burying in an approved landfill.

## MATERIAL SAFETY DATA SHEET

Additional information is listed on the Material Safety Data Sheet and is available from Ensystex on request. Or visit our web site at [www.Ensystex.co.nz](http://www.Ensystex.co.nz).

## NOTICE

Ensystex New Zealand Ltd warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions for use and under normal conditions of use. No warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label or under abnormal conditions.



Batch No.:

Date of Manufacture:

**L-NZ-RDTB1018 2.02 10.21**